

TRITC Antibody Labeling Kit

T1491948

Storage For detailed storage information, please refer to the kit contents.

Introduction

The TRITC Antibody Labeling Kit provides all necessary reagents for antibody labeling, including Tetramethylrhodamine (TRITC), buffers, and a spin desalting column. The reactive dye efficiently conjugates to lysine residues in antibodies or proteins, forming a stable covalent product. The included spin purification column enables rapid removal of excess unreacted dye, with a recovery rate of 70%–95%. The kit contains 5 vials of reactive dye. Each vial is designed for labeling 100 µg of antibody.




Fig. 1 TRITC Antibody Labeling Kit (T1491948) Labeling Principle Schematic

Kit Contents

T1491948	Component	5 reactions	Storage	Quantity Per reaction
T1491948A	TRITC	5 vials	-20°C. Store in the dark.	1 vial for labeling 100 µg of antibody
T1491948B	Sodium Carbonate-Bicarbonate	1 vial	RT.	Prepare according to instructions
T1491948C	Spin Desalting Column	5 EA	2-8°C. Do not freeze.	1 EA for 1 reaction
T1491948D	Collection Tubes	10 EA	RT.	2 EA for 1 reaction

Required materials not supplied

1. Microcentrifuge capable of 1,000 × g.
2. Desired antibody for labeling (free of BSA or any carrier protein).
3. PBS buffer (pH 7.2-7.4).

Matters needing attention

1. The purified antibody should be in a buffer that does not contain primary amines (for example, ammonium ions, Tris, glycine, ethanolamine, triethylamine, glutathione) or imidazole. All of these substances significantly inhibit protein labeling.
2. Impure antibodies or antibodies stabilized with bovine serum albumin (BSA) or gelatin will not be labeled well.
3. Concentrate the antibody to ≥ 1 mg/mL.
4. Do not reuse the purification resin.
5. Before opening the vials, warm all components to room temperature.
6. The reactive dyes should be protected from prolonged exposure to light.

Instructions for Use

1. Prepare the reagents
 - (1) Prepare a 1 M Sodium Carbonate-Bicarbonate Buffer: Add 1 mL of ultrapure water to the tube containing sodium carbonate-sodium bicarbonate powder. Cap the tube and shake until the powder is completely dissolved to obtain 1 M sodium carbonate-bicarbonate buffer (pH 9.2).

Note: The prepared 1M Buffer is stable for 1-2 weeks when stored at 2-8°C.

2. Label the antibody
 - (1) If the antibody to be labeled has a concentration of ≥ 1 mg/mL and is in an appropriate buffer, dilute it to 1 mg/mL, and add a 10% volume of 1 M Carbonate-Bicarbonate Buffer. If the antibody is a powder lyophilized from an appropriate buffer, prepare a 1 mg/mL solution of the antibody by adding an appropriate amount of 0.1 M Carbonate-Bicarbonate Buffer to the antibody. Prepare 0.1 M Carbonate-Bicarbonate Buffer by diluting the 1 M solution 10-fold with ultrapure water.
 - (2) Transfer 100 μ L of the antibody solution to the vial of reactive dye. Cap the vial and gently invert it a few times to fully dissolve the dye.

Note: To visually ensure that the dye has fully dissolved, peel the label off the vial of reactive dye.

- (3) Incubate the solution for 2 hours at room temperature in the dark. The reaction can be carried out on a shaker or mixer, recommended speed for flipping up and down is 25rpm. If a mixing instrument is not used during the reaction process, the reaction solution should be mixed upside down every 10 minutes.

Note: During the incubation period, proceed to steps 3 below, to prepare a spin column for the purification of the labeled protein.

3. Prepare the spin column
 - (1) Loosen the cap on a spin column, twist the tab off of the bottom, then place the column into a collection tube.
 - (2) Centrifuge the column-tube assembly at $1,000 \times g$ for 2 minutes to remove the storage solution.

Note: When using a fixed-angle rotor, place a mark on the side of the column that faces away from the rotor center. For all subsequent centrifugation steps, place the column in the microcentrifuge with the mark facing away from the rotor center.

- (3) Discard the flowthrough, then place the column back into the collection tube.
- (4) Add 500 μ L of PBS Buffer, then centrifuge the column-tube assembly at $1,000 \times g$ for 2 minutes to equilibrate the column. Repeat this step (3) 3 times, discarding the buffer from the collection tube each time.

4. Purification with Desalting Column

- (1) Transfer the equilibrated column into a new collection tube.
- (2) Carefully pipette the entire reaction mixture onto the center of the column.
- (3) Centrifuge the column-tube assembly at $1,000 \times g$ for 2 minutes. The purified antibody conjugate is in the collection tube.

5. Determine the antibody concentration and DOL (Optional)

- (1) Use the absorption spectrophotometer to measure the absorption spectrum of the purified label, and record the absorbance values at 280 nm and 557nm (with a 1-cm path length).
- (2) The following formula can be used to calculate the antibody concentration:

$$\text{Antibody concentration (mg/mL)} = (A_{280} - CF_{280} \times A_{557}) \times 150000 / \epsilon_{\text{IgG}}$$

- (3) The following formula can be used to calculate the degree of labeling:

$$\text{DOL} = (A_{498} / \epsilon_{\text{TRITC}}) / [(A_{280} - CF_{280} \times A_{557}) / 210,000]$$

Parameter	Meaning	Value
DOL	Represents the molar ratio of the fluorescent dye to the target molecule (e.g., antibody or protein) in the conjugate	-
A_{557}	Absorbance value of TRITC at a wavelength of 557 nm	Measured value
ϵ_{TRITC}	Molar extinction coefficient of TRITC ($\text{M}^{-1}\text{cm}^{-1}$)	100000
A_{280}	Absorbance value of the TRITC-labeled antibody at a wavelength of 280 nm	Measured value
CF_{280}	Correction factor for the absorbance value of TRITC at 280 nm	0.34
ϵ_{IgG}	Molar extinction coefficient of the IgG antibody at 280 nm ($\text{M}^{-1}\text{cm}^{-1}$)	210000

6. Storage

- (1) Add 0.05–0.2% Proclin 300 or 0.05% sodium azide, along with a protein stabilizer (such as 0.1% BSA), to the labeled protein. Store protected from light at $2-8^{\circ}\text{C}$ for stable preservation up to six months. Alternatively, add an equal volume of glycerol and store at -20°C for stable preservation up to six months.