

Rifamycin sodium enzyme

rp216614

Store at 2-8°C

Introduction:

Rifamycin sodium enzyme is a water-soluble protein using recombinant genes, presenting as a dark brown liquid. It is a recombinant gene protein constructed according to the chemical structure of rifamycin sodium. The active center of its gene fragment can disrupt the chemical structure of rifamycin sodium, causing the drug to lose its antibacterial properties after ring opening or chain breaking.

Notes:

1. When conducting the sterility test of antibiotics, use a manual syringe to draw sterile water and inject it into the vial of the drug. Shake it well to dissolve the drug, and then draw out the solution. Add it to a 500 ml solution of 0.9% sodium chloride and shake it well. Do not use the needle on the incubator to dissolve and inject the solution. Avoid high-concentration solutions passing through the filter membrane, which may make it difficult to rinse thoroughly.
2. Here, it is emphasized again: when conducting the sterility test of antibiotics, be sure to use a manual syringe to inject sterile water to dissolve the sample, and then transfer the dissolved sample to a 500 ml solution of 0.9% sodium chloride and shake it well, so as to prevent the sample from having a too high local concentration and causing the situation where it is difficult to rinse thoroughly when passing through the filter membrane.
3. During the sterility test, take 3 ml of the enzyme and add 2 ml of sterile water, and shake it well to make a diluted solution of rifamycin sodium enzyme. Take 2 ml of the diluted enzyme solution and put it into 1500 ml of the rinsing solution and shake it well. After the rinsing solution has finished rinsing the filter membrane of the incubator, pump out all the solution. Use a manual syringe to pierce into the breathing ports of the three incubators, and add 1 ml of the diluted enzyme to each incubator. Try to make the enzyme spread evenly over the entire surface of the filter membrane. Let it stand for 10 minutes to ensure that the high-concentration enzyme fully contacts the filter membrane of the incubator, so as to destroy (neutralize, inactivate) the remaining rifamycin sodium drugs on the filter membrane. Then pump in the corresponding culture medium and shake it well. Add 1 ml of the corresponding test bacteria to the positive control tube.
4. Adding 2 ml of the diluted solution of rifamycin sodium enzyme to the rinsing solution can remove the antibacterial properties of the small amount of rifamycin sodium remaining inside the filter membrane.

5. Adding 1 ml of the diluted solution of rifamycin sodium enzyme to each of the three incubators can remove the antibacterial properties of the small amount of rifamycin sodium remaining on the inner wall of the incubator and the surface of the filter membrane.
6. Gently shake the positive control tube once in the morning and once in the afternoon every day.
7. Customers can conduct methodological verification according to the above methods, or conduct verification according to the actual operation situation.

